

Resolution Analytical Column



TRINITY BIOTECH
KANSAS CITY, MO 64132 USA
www.trinitybiotech.com







15-090-3 2018-10-28

01-05-0015

CONT

1 Each



Trinity Blotech plc Brey, Co. Wickiow, Ireland Tek +353 1 276 9860 Fax: +353 1 276 9888

Certificate of Analysis

Intended Use The Trinity Biotech Resolution Analytical Column is intended for use with the Ultra2 Resolution Variants Analyzer. No substitutions are permitted, registered, cleared or authorized. No other uses are intended, registered, cleared or authorized.

The Ultra2 Resolution Variants Analyzer is intended for the qualitative detection of hemoglobin variants and quantitative measurement of hemoglobin HbF and HbA2 in human capillary, venous whole blood and dried blood spots. HbF and HbA2 are used for evaluating samples for thalassemia and other hemoglobinopathies. For in vitro diagnostic use only.

Performance Analysis

BASELINE ACCEPTABILITY

Standard Result Initial baseline flat with no deflection higher than 20 mm above starting base.

The initial baseline is flat with no deflection on the printed chromatogram greater than 20mm above the starting base.

HbA2 INTEGRATION ACCEPTABILITY

Standard Result Baseline separation of A2 from HbA and HbS with symmetrical peak shape. HbA2 is symmetrical and has baseline separation from HbA and HbS.

A2,F CONTROL RESULTS

Standard Result HbF and HbA2 recovery within limits for normal and abnormal controls. HbF and HbA2 values are within specs stated in the package insert.

Standard

FASC RETENTION TIME (QUICK SCAN) - HbF PEAK

Standard HbF elutes between 1.4 and 1.6 minutes.

Result The retention time of HbF is between 1.4 and 1.6 minutes.

Standard

FASC RETENTION TIME(QUICK SCAN) — HbA PEAK HbA elutes between 2.2 and 2.6 minutes.

Result

The retention time of HbA is between 2.2 and 2.6 minutes.

Standard

FASC RETENTION TIME (QUICK SCAN) – HbS PEAK
HbS elutes between 2.8 and 3.2 minutes.

Result

The retention time of HbS is between 2.8 and 3.2 minutes.

Standard Result FASC RETENTION TIME (QUICK SCAN) – HbC PEAK HbC elutes between 3.3 and 3.6 minutes and returns to baseline. The retention time of HbC is between 3.3 and 3.6 minutes.

CROSS COLUMN PRESSURE

Standard Result The cross-column pressure is less than 75bar at 1.5mL/minute. At 1.5mL/min the cross-column pressure is less than 75bar.

AUTHORIZED REPRESENTATIVE APPROVAL

Date

11/17/2016

Quality Control

Resolution Analytical Column

REF

01-05-0015

1 each

Pour d'autres langues Für andere Sprachen Para otras lenguas Per le altre lingue

Dla innych języków

Para outras linguas Για τις άλλες λώσσες För andra språk For andre språk



www.trinitybiotech.com

INTENDED USE

The Trinity Biotech Resolution Analytical Column is intended for use with the Ultra2 Resolution Variants Analyzer. No substitutions are permitted, registered, cleared or authorized. No other uses are intended, registered, cleared or authorized.

The Ultra2 Resolution Variants Analyzer is intended for the qualitative detection of hemoglobin variants and quantitative measurement of hemoglobin HbF and HbA2 in human capillary, venous whole blood and dried blood spots. HbF and HbA2 are used for evaluating samples for other hemoglobinopathies. thalassemia For in vitro diagnostic use only. IVD

SUMMARY AND EXPLANATION OF TEST

Utilizing a High Performance Liquid Chromatography (HPLC) system hemoglobins are separated, quantified and presumptively identified by comparison to reference peaks of Hb F, A, S and C. The ion-exchange column that has been equilibrated with respect to pH and ionic strength is used to separate the hemoglobin species.

Resolution Analytical Columns are performance validated to assure accuracy and precision with our assays and systems for the measurement of hemoglobin species.

The Column is ready for use

STORAGE AND STABILITY



Store at 2 – 8°C for long term storage.

Columns that are refrigerated at 2-8°C are stable for two years when kept tightly closed. Columns that are placed into service have a limited shelf life of a few weeks and will be gradually consumed once placed into service.



EXP See the column label for the expiration date. DO NOT USE after the expiration date.



Allow the column to warm to room temperature before installing on the instrument, Equilibrate the column at 1.5mL/min at the Standby conditions for 15 minutes before use.

PRECAUTIONS

For in vitro diagnostic use only. Avoid skin contact. Consult the product MSDS for safety information. This column is used in conjunction with blood testing equipment and warrants handling under universal precaution procedures for safety.

ORDERING INFORMATION			
	REF	ltem	Quantity
01	-05-0015	Resolution Analytical Column	1 each

COLUMN LIFE

Column life will vary depending on diligence in system maintenance (regular and preventative


maintenance, as scheduled and using manufacturer specified items). Column life will vary depending on weekly test throughput (low throughput and infrequently used systems may not achieve the average number injections.) Column life will vary depending on diligence in column maintenance (trit changes, reversing column direction (flipping), proper shutdowns (nightly/weekends) to preserve the column. Column life will vary depending on diligence in reagent management (closed containers, no topping-off, replacement of fouled check-valves if reagent is allowed to run dry). Column life will vary depending on diligence in control management (careful preparation according to PI reconstitution instructions, careful preservation according to PI instructions).

Any series of columns experiencing reduced life on the same instrument is an indication of a system or operation issue (or very low weekly test throughput). Systems in need of routine or preventive maintenance will experience reduced column life. For these systems, although changing the column provides improvement, it is not the underlying cause, and short column life will continue until the issue is properly addressed.

NOTE: Column warranty claims must include the following supporting information: maintenance schedule (date of last PM), column change report (or monthly inventory) report, chromatography (including cover page and header information), the number of injections, and any follow-up information requests made. Any claim with missing information, as specified above, cannot be processed.

USED COLUMN RETURN

Please return used columns each month. Columns are tracked by serial number. Return columns may be tested and reviewed for the purposes of product improvement.





Trinity Biotech Kansas City, MO 64132 Tel. 1 800-325-3424 Fax: 1816-361-1974



Trinity Biotech plc Bray Co. Wicklow, Ireland Tel, 353 1 2769800 Fax 353 1 2769888 www.trinitybiotech.com



w 11/17/20/6 MSL 11-1716